

Chandra and *Swift* follow-up observations of the intermediate mass black hole in ESO243-49

N. A. Webb, D. Barret, O. Godet

Université de Toulouse, UPS, CESR, 9 Avenue du Colonel Roche, F-31028 Toulouse Cedex 9, France

`natalie.webb@cesr.fr`

M. Servillat

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, MS-67, Cambridge, MA 02138, USA

S. A. Farrell

Department of Physics and Astronomy, University of Leicester, University Road, Leicester, LE1 7RH, UK

S. R. Oates

Mullard Space Science Laboratory/UCL, Holmbury St. Mary, Dorking, Surrey, RH5 6NT, UK

ABSTRACT

The brightest Ultra-Luminous X-ray source HLX-1 in the galaxy ESO 243-49 provides strong evidence for the existence of intermediate mass black holes. As the luminosity and thus the mass estimate depend on the association of HLX-1 with ESO 243-49, it is essential to confirm its affiliation. This requires follow-up investigations at wavelengths other than X-rays, which in-turn needs an improved source position. To further reinforce the intermediate mass black hole identification, it is necessary to determine HLX-1's environment to establish whether it could potentially form and nourish a black hole at the luminosities observed. Using the High Resolution Camera onboard *Chandra*, we determine a source position of $RA=01^h10^m28^s.3$ and $Dec=-46^\circ04'22''.3$. A conservative 95% error of $0.3''$ was found following a boresight correction by cross-matching the positions of 3 X-ray sources in the field with the 2MASS catalog. Combining all *Swift* UV/Optical Telescope *uvw2* images, we failed to detect a UV source at the *Chandra* position down to a 3σ limiting magnitude of 20.25 mag. However, there is evidence that

the UV emission is elongated in the direction of HLX-1. This is supported by archival data from *GALEX* and suggests that the far-UV emission is stronger than the near-UV. This could imply that HLX-1 may be situated near the edge of a star forming region. Using the latest X-ray observations we deduce the mass accretion rate of a $500\text{ M}_{\odot}$ black hole with the observed luminosity and show that this is compatible with such an environment.

Subject headings: X-rays: individual — X-rays: binaries — accretion, accretion disks

1. Introduction

The brightest Hyper Luminous X-ray source (HLX-1) was discovered serendipitously with *XMM-Newton* on 2004 November 23 (Farrell et al. 2009) in the outskirts of the edge-on spiral galaxy ESO 243–49, at a redshift of 0.0224 (Afonso et al. 2005). Its $0.2 - 10\text{ keV}$ unabsorbed X-ray luminosity, assuming isotropic emission and the galaxy distance (95 Mpc), exceeded $1.1 \times 10^{42}\text{ ergs s}^{-1}$. Follow-up observations with *XMM-Newton* and the *Swift* X-ray telescope (XRT) revealed that the source was variable in X-rays by more than one order of magnitude, and that luminosity changes were accompanied by changes in the spectral shape, in a way similar to Galactic black hole systems (Godet et al. 2009). As argued before, either super-Eddington accretion or beaming of the X-ray emission could account for X-ray luminosities up to $\sim 10^{40}\text{ ergs s}^{-1}$ for stellar mass black holes ($10\text{--}50\text{ M}_{\odot}$), but would require extreme tuning to explain an X-ray luminosity of $10^{42}\text{ ergs s}^{-1}$. Hence, HLX-1 is an excellent intermediate mass black hole (IMBH) candidate (Farrell et al. 2009).

To verify the IMBH nature it is essential that its association with the host galaxy is confirmed. To do this, an improved position is crucial in order to carry out multi-wavelength follow-up observations. Subsequently, it is necessary to determine whether the source resides in an environment which can provide sufficient material to power such a massive black hole (Milosavljević et al. 2009). Again multi-wavelength observations are the key to determining whether this object is in a star cluster, a star forming region or a globular cluster. As ULXs emit mostly in the X-ray domain, X-ray observations are thus the key to understanding how much material is required to feed the black hole as the luminosity is directly related to the mass accretion rate. Therefore constant monitoring of the X-ray emission will allow us to place constraints on the quantity of material necessary in the neighbouring environment.

Here we present *Chandra* observations of HLX-1 with the High Resolution Camera (HRC) accorded under the Directors Discretionary Time (DDT) program, which has allowed

us to revise and improve the position of HLX-1, along with *Swift* XRT data revealing that the X-ray luminosity of this source remains above 10^{42} ergs s $^{-1}$, following its recent rebrightening in August 2009 (Godet et al. 2009). We discuss the mass accretion rate of HLX-1 using these observations. We also reveal evidence for the nature of the environment around HLX-1 through ultra-violet imaging with the *Swift* UV/Optical Telescope (UVOT); a finding independently supported by archival *Galaxy Evolution Explorer* (GALEX) observations.

2. Data analysis

2.1. The first HRC-I observation

We obtained a 1 ks DDT observation of HLX-1 with the HRC-I camera onboard *Chandra* on 2009 July 4 (ObsID: 10919). We extracted all the events in the detector (i.e. the background) and plotted the number of counts per energy channel (PI) with the CIAO v4.1.1 task WAVDETECT. Between PI = 25 and 120 the background is lower and the sensitivity of the instrument is higher (see e.g. Cameron et al. 2007). We thus filtered the event list to include only these energies and performed source detection with WAVDETECT utilising an exposure map. No source is detected within the *XMM-Newton* error circle of HLX-1, and only 2 sources are detected in the field of view. The two sources that are detected correspond to known *XMM-Newton* sources: 2XMM J011050.4-460013 which was similar in flux to the *XMM-Newton* detection of HLX-1, and 2XMM J010953.9-455538. These sources have respectively 17 and 6 counts in the HRC-I image, which gives 0.018 and 0.006 ct s $^{-1}$. Therefore, HLX-1 must have dropped in flux down to at most 0.006 ct s $^{-1}$ (the faintest source detected), compared to 0.03 ct s $^{-1}$ expected from the flux in the most recent *XMM-Newton* observation. This faint flux is confirmed using *Swift* observations taken one month later (Godet et al. 2009).

2.2. The second HRC-I observation

Following the re-brightening of HLX-1 (Godet et al. 2009) we obtained a second deeper DDT observation of 10 ks with the *Chandra* HRC-I 2009 on August 17 (ObsID: 11803). We generated several images of the HRC-I field of view with a binning of 4, 8, 16 and 32 pixels, and performed source detection using WAVDETECT. A total of 11 sources were detected, including a source consistent with the position of HLX-1 with a net count rate of 0.098 ± 0.003 ct s $^{-1}$. We cross-matched the entire source list with the 2MASS catalogue (Skrutskie et al. 2006) and found 3 matching objects which appear to be the real counterparts

to the X-ray sources. A boresight correction was applied to the X-ray image which lead to a small shift of $0.285''$. After applying the astrometry correction to the image `WAVDETECT` was run again. The final corrected position for HLX-1 was found to be $RA = 01^h10^m28^s.3$ and $Dec = -46^\circ04'22''.3$, which is consistent with the previous derived positions. The error on this position was then estimated from the 1σ error of the 2MASS catalogue at that position ($0.1''$ see Skrutskie et al. 2006), the 1σ `WAVDETECT` 2D error ($0.030''$), and the root mean square (RMS) of the alignment of the X-ray image on the 2MASS reference stars ($0.003''$). Combining these errors, a total 95% error of $0.3''$ was derived.

2.3. The *Swift* UVOT data

The *Swift* UVOT observed the field of ESO 243-49 on 2009 August 5, 6, 16, 18, 19 and 20 for a total exposure of 38 ks. Observations were performed in the *uvw2* (~ 1600 - $2500\text{\AA}$) filter only. Data reduction and analysis of the data were done using the software release HEADAS 6.6.2 and version 20090630 of the UVOT calibration files. At the location of the core of ESO 243-49 there is an extended object (Figure 2), with some hints of an elongated emission towards the position of HLX-1. No other source is observed above the flux level of this galaxy at the *Chandra* position of HLX-1.

We cannot determine an upper limit from subtracting the galaxy contribution at that position as we do not know when or even if HLX-1 stops emitting in the *uvw2* wavelength range. A source at the X-ray position would only be discernible from the galaxy if the source is significantly brighter than the emission at that position. Therefore, to determine an upper limit to the detection of the ULX above the flux of the galaxy, we first co-added the individual observations. We then placed 7 circular apertures of $3''$ radius on the image, one at the location of the X-ray source and 6 others around an ellipse on the galaxy of similar galaxy isophotal brightness to the region containing the source. Then using a background region of $20''$ located in a source free region close to the galaxy, we determined the magnitude of the galaxy in each aperture using the *Swift* tool `UVOTSOURCE`. To be compatible with the calibration, which is determined for a $5''$ aperture (Poole et al. 2008), we applied an aperture correction to the count rate using a table of aperture correction factors contained within the calibration files. We determined the standard deviation of the magnitudes determined from the 7 apertures to be 0.10 mag. We then took the 3σ upper-limit for detecting a source at the *Chandra* position to be the magnitude determined from the source region located at the *Chandra* position minus 3 times the standard deviation, which gives a 3σ upper-limit of 20.25 mag.

2.4. The *GALEX* data

Given the possible detection of extended emission in the *uvw2*, we retrieved archived data taken by *GALEX* in the near- (NUV, $\sim 1800\text{--}2800\text{\AA}$) and far-UV (FUV, $\sim 1500\text{--}2000\text{\AA}$) bands. The field of HLX-1 was observed by *GALEX* as part of the deep survey on 2004 September 27 for ~ 13 ks in the NUV and ~ 8 ks in the FUV. A clear extension towards the location of HLX-1 from the bulge of ESO 243-49 can be seen in both images (see Figure 3), with the dominant emission occurring in the FUV. No point source was detected coincident with the position of HLX-1 in either band. In order to determine magnitude upper limits, we measured the counts within regions centered on the *Chandra* position with radii corresponding to the 80% encircled energy radii ($3''$ and $4''$ for the NUV and FUV respectively). The total count rate inside six similar sized regions located around the galaxy, at the same isophotal brightness as the location of HLX-1 were also determined. Using a source-free area of the image near the galaxy with radii of $6''$ and $8''$ for the NUV and FUV respectively we measured the total background rates. These were scaled to the same extraction region size as used for the source regions. All the count rates (both source and background) were then corrected for the wings of the PSF outside the extraction region. Net count rates were calculated by subtracting the background rates from the source rates. Magnitudes were then calculated for the net rates using the zero points given in Morrissey et al. (2007). The standard deviation for the magnitudes from each of the 7 regions were 0.3 and 0.2 mag for the NUV and FUV respectively. We then determined the 3σ upper limits as the magnitudes derived for the region centered on the HLX-1 *Chandra* position minus 3 times the standard deviation, in the same way as for the UVOT data. This gives upper limits of 20.4 mag and 21.4 mag for the NUV and FUV respectively.

2.5. The XRT data

The third Target-of-Opportunity (ToO) observation of our *Swift*-XRT monitoring campaign was performed on 2009 October 02 (hereafter designated S4 as it is the fourth of our *Swift* observations of this source, see Godet et al. 2009) for 9.4 ks. All the *Swift*-XRT Photon Counting data were processed using the tool `XRTPIPELINE v0.12.3`. To analyse the data, we used the same method as described in Godet et al. (2009). This new observation revealed that the HLX-1 0.3–10 keV count rate dropped from $\sim 3.3 \times 10^{-2}$ to $\sim 1.9 \times 10^{-2}$ count s^{-1} when compared to the previous observation taken on 2009 August 16 (S3; see Godet et al. 2009). Along with this reduction in count rate, we also observe a spectral softening as illustrated in Fig. 1. The hardness, defined as the ratio of the 0.3–1 keV count rate over the 1–10 keV count rate, is 0.26 ± 0.05 in S4, compared to a value of 0.42 ± 0.04 in S3. The

errors are quoted at the 1σ level. Both a unique absorbed disk black-body (DBB) and an absorbed power-law (PL) model give comparable fits ($\chi^2/dof = 9.8/6$ and $\chi^2/dof = 10/6$, respectively). The n_H -value was fixed at $4 \times 10^{20} \text{ cm}^{-2}$ (Farrell et al. 2009) as the poor quality of the spectra meant that we were not able to derive a meaningful value for this quantity, although the allowed values always included the fixed one. For the DBB model, we obtained a temperature of $kT = 0.20_{-0.02}^{+0.03} \text{ keV}$, consistent with the disc blackbody temperature found in previous observations with *XMM-Newton* (see Farrell et al. 2009) and a 0.2–10 keV unabsorbed luminosity of $1.0 \pm 0.3 \times 10^{42} \text{ erg cm}^{-2} \text{ s}^{-1}$, while for the PL model we obtained a value for the photon index of $\Gamma = 3.5 \pm 0.3$ and a 0.2–10 keV unabsorbed luminosity of $1.9_{-0.4}^{+0.5} \times 10^{42} \text{ erg cm}^{-2} \text{ s}^{-1}$. A full discussion of the spectral nature of HLX-1 can be found in Godet et al. (2009). We add here that the spectral state observed for S4 is intermediate between S1 and S3, which appears to portray smooth spectral evolution that resembles that of black hole binaries, supporting the interpretation in Godet et al. (2009). However, here we exploit only the luminosity so as to constrain mass accretion limits, as discussed below.

3. Discussion and conclusions

Using *Chandra* observations of HLX-1, we have determined an improved position with a 95% confidence error radius of $0.3''$. This is sufficiently small that we should eventually be able to identify the source at other wavelengths, and eventually perform broad band spectroscopic observations.

Although no point-like source is detected with UV observations at this position, HLX-1 appears to be situated near the edge of a region of UV excess stretching from the nucleus of ESO 243-49 towards the *Chandra* position. However, it is not yet certain that this emission is related to HLX-1, but the fact that no similar extended emission is seen in the radio, infra-red, optical, or X-ray domains (Farrell et al. 2009, Webb et al. in preparation), makes it unlikely that this is either a foreground or background source. Follow-up observations with a higher resolution instrument would also help to confirm the association and the extended nature, as the low resolution of *GALEX* ($5''$ Full Width Half Maximum (FWHM)) and the *UVOT* (better at $2.9''$ FWHM) would mean that it would be difficult to resolve a source at the centre of the galaxy and a second towards the position of HLX-1. However, the fact that this emission appears to be stronger in the FUV could hint towards star formation taking place in that region, as the UV flux primarily originates from the photospheres of O- through later-type B-stars ($M > 3M_{\odot}$), and thus measures star formation averaged over a $\sim 10^8 \text{ yr}$ timescale (e.g. Lee et al. 2009). Starburst environments are thought to be able to generate IMBHs through runaway collisions and mergers of massive




Fig. 1.— *Swift*-XRT PC grade 0–12 unfolded spectra of HLX-1: S4 = 2009-10-02 (black) and S3 = 2009-08-16 (red). The solid lines correspond to the best-fit models. For S4, we used a unique absorbed disk black body. The n_H -value was fixed at $4 \times 10^{20} \text{ cm}^{-2}$ in both cases.

stars (Freitag, Rasio & Baumgardt 2006). Further, the massive stars present in such environments can supply the necessary material to be accreted onto the black hole to provide the luminosities observed if one is captured by the black hole and then proceeds to main sequence Roche-lobe overflow (Hopman, Portegies Zwart & Alexander 2004). An alternative situation is described by Sun et al. (2010) where a trail of star formation could be created by ram-pressure stripping of gas and stars from a dwarf galaxy which has recently interacted with ESO 243-49. In this case HLX-1 could have been an intermediate mass black hole which was once at the centre of the dwarf galaxy, but has now had most of the gas and stars stripped from it via the gravitational interaction with ESO 243-49. This may resemble a globular cluster if detected in the optical domain.




Fig. 2.— *Swift*-UVOT un-smoothed *uvw2* image of ESO 243-49. The white contours show the orientation of the galaxy in the J-band (Webb et al. in preparation). The black circle indicated by the white arrow is centered on the *Chandra* position of HLX-1, with the radius representing the 95% error bounds.

To determine whether accretion from the medium around the stripped galaxy is possible, we use the latest X-ray observation of HLX-1, which has a 0.2–10 keV unabsorbed luminosity of $1.9^{+0.5}_{-0.4} \times 10^{42}$ erg cm $^{-2}$ s $^{-1}$ if we fit with a power law model. This is the highest luminosity that we have observed over the last five years whilst this source has been bright (previous observations with *Rosat* in the early nineties gave non-detections, confirming that the source was more than a factor 10 fainter than in these observations, Webb et al. in preparation). HLX-1 is not bright in the radio, infra-red, optical or NUV domains (Farrell et al. 2009, Webb et al. in preparation), so we can assume that the majority of the emission is in the X-rays and we can use this luminosity (L) to deduce the approximate mass accretion rate ($\dot{M}$), where $L = \eta \dot{M} c^2$ and $\eta = GM/Rc^2$ (M is the mass of the black hole and R the innermost stable circular orbit i.e. 6 times the Schwarzschild radius, therefore supposing a non-rotating black hole). We assume a mass of $500 M_{\odot}$. We find $\dot{M} \sim 2.1 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$.




Fig. 3.— Archival *GALEX* near-UV (left) and far-UV (right) images of ESO 243-49. Both images have been smoothed using a 2 pixel radius Gaussian smoothing function. The white contours show the orientation of the galaxy in the J-band (Webb et al. in preparation). The black circles indicated by the white arrows are the *Chandra* position of HLX-1, with the radii indicating the 95% positional error of $0.3''$.

This is well within the ranges of the matter available for accretion predicted by Sun et al. (2010), therefore supporting the idea that HLX-1 *could* be embedded in a star forming region and that it may have originated from a stripped dwarf galaxy.

We thank Neil Gehrels and the *Swift* team for according us the *Swift* observations as well as Harvey Tananbaum and the *Chandra* team for approving the two *Chandra* DDT observations. S.A.F. and S.O. acknowledge STFC funding. This research has made use of data obtained from the *Chandra* Data Archive and software provided by the *Chandra* X-ray Center. We thank the *GALEX* collaboration and the Space Telescope Science Institute for providing access to the UV images used in this work. *GALEX* is a NASA Small Explorer Class mission. We thank Simon Rosen for valuable discussions and we are grateful to the referee for comments which improved this paper.

Facilities: *Chandra*, *Swift*, *GALEX*

REFERENCES

- Afonso, J. et al. 2005 ApJ, 624, 135
- Cameron, P. B., Rutledge, R. E., Camilo, F., Bildsten, L., Ransom, S. M., & Kulkarni, S. R. 2007, ApJ, 660, 587
- Farrell, S. A., Webb, N. A., Barret, D., Godet, O. & Rodrigues, J. M. 2009, Nature, 460, 73
- Freitag, M., Rasio, F. A., & Baumgardt, H., 2006, MNRAS, 368, 121
- Godet, O., Barret, D., Webb, N. A., Farrell, S. A., & Gehrels, N. 2009, ApJL, 705, 109
- Hopman, C., Portegies Zwart, S. F., & Alexander, T., 2004, ApJ, 604, 101
- Lee, J.C., de Paz, A. G., Temonti, C., & et al., 2009, ApJ, 706, 599
- Miller, M. C., & Colbert, E. J. M. 2004, International Journal of Modern Physics D, 13, 1
- Milosavljević, M., Couch, S. M., & Bromm, V. 2009, ApJL, 696, L146
- Morrissey, P., et al. 2007, ApJSS, 173, 682
- Poole, T. S., et al. 2008, MNRAS, 383, 627
- Skrutskie, M. F., et al. 2006, AJ, 131, 1163
- Sun, M., Donahue, M., Roediger, E., & et al. 2010, ApJ, 708, 946